Checking your blood sugar

How frequent blood sugar checks can help you

Checking your blood sugar yourself is an important part of managing diabetes. Checking often will tell you:

- If your insulin or other diabetes medicine is working
- How physical activity and the foods you eat affect your blood sugar

You'll usually feel better and have more energy when your blood sugar stays at or near normal. Managing your blood sugar can also reduce your risk of developing problems from diabetes.



How to check your blood sugar

You can check your own blood sugar using a meter. Many different kinds of blood sugar meters are available today. Your diabetes care team can help you choose one and show you how to use it.

When to check your blood sugar

You and your diabetes care team will decide when and how often you will check your blood sugar. The table below shows some times when you might want to check and why.

When your team may want you to check	Why you should check			
When you wake up	To see if your blood sugar is staying under control while you're asleep (called fasting blood sugar)			
Before meals	To know what your blood sugar is before you eat and to know if you will need to adjust your mealtime (prandial) insulin			
1 or 2 hours after you start your meal	To see how the food you eat and insulin dose affect your blood sugar			
Before, during, and after physical activity (depending on the length of time of your activity)	To see how being active affects your blood sugar			
At bedtime	Depending on the medicine that you take			

Keeping a blood sugar tracker

It's important to write down your blood sugar levels so that you can keep track of what makes them go up or down.

See the last page for a tracker that you can use to record your blood sugar and write down other important information for your diabetes care team.



Checking your blood sugar

Time	Goals for many adults with diabetes	Your goals			
Before meals	80 to 130 mg/dL				
1 to 2 hours after the start of a meal	Less than 180 mg/dL				
A1C	Less than 7%				
American Diabetes Association. Standards of medical care in diabetes—2016. <i>Diabetes Care</i> , 2016;39(suppl 1):S1-S112.					

Setting your blood sugar goals

The table above lists blood sugar goals for many adults with diabetes. You and your diabetes care team will set the goals that are right for you. Write your goals in the last column.

Knowing your A1C

The A1C test measures your estimated average blood sugar level over the past 2 to 3 months. It's like a "memory" of your blood sugar levels. It shows how well you're controlling your blood sugar levels over time. Your A1C and your blood sugar levels go up and down together. The table to the right shows how A1C relates to the estimated average blood sugar level.

Lowering your A1C to below 7% reduces your risk of problems from diabetes. Therefore, the A1C goal for most people is less than 7%.



It is recommended that you get an A1C test:

- At least 2 times a year if your blood sugar is under good control
- 4 times a year if you are not meeting your goals or if your treatment has changed

A1C level	Average blood sugar				
6%	126 mg/dL				
7%	154 mg/dL				
8%	183 mg/dL				
9%	212 mg/dL				
10%	240 mg/dL				
11%	269 mg/dL				
12%	298 mg/dL				
American Diabetes Association. Standards of medical care in diabetes—2016. Diabetes Care. 2016;39(suppl 1):S1-S112.					

If you have any questions, be sure to talk with your diabetes care team. They are there to help!

For more information, visit Cornerstones4Care.com

Novo Nordisk Inc. grants permission to reproduce this piece for nonprofit educational purposes only on condition that the piece is maintained in its original format and that the copyright notice is displayed. Novo Nordisk Inc. reserves the right to revoke this permission at any time.

Cornerstones4Care® is a registered trademark of Novo Nordisk A/S



Checking your blood sugar

Cornerstones4Care

Your blood sugar tracker

Diabetes changes over time. That is why your treatment may also need to change over time. For example, your doctor may tell you to add a basal insulin to your diabetes pills. Or if you already take a basal insulin, your doctor may tell you to add mealtime insulin.

The blood sugar tracker on the next page can help you keep track of your diabetes medicines and any changes in dosage and timing that your doctor may tell you to make. It can also help you keep track of how much mealtime insulin you took and when you took it.

Your diabetes care team can provide you with a blood sugar tracker and show you how to use it. Or call **1-800-727-6500** to have a tracker sent to you.

Carb intake ▶

Mealtime insulin dose ► ☐ Breakfast_

Other >

20 carba


BP: 120/80

EXAMPLE

Here's how to use the blood sugar tracker

- 1 Write down the date for the start of the week. (You can start tracking on any day of the week)
- 2 Write the name(s) and dose(s) of your diabetes medicine(s)
- Write the time and your blood sugar readings in the "before" and "after" spaces. After-meal readings are usually taken 1 to 2 hours after you start your meal. Nighttime readings may be taken during the night as needed
- 4 If you are counting carbs, write how many grams of carbs you ate
- If your doctor has told you to use mealtime insulin when you eat, write how many units you took
- 6 Add notes on anything else you might want to track (such as blood pressure or weight)

Dinner 2 units



Lunch___

Your blood sugar tracker

Date:	1		/
Date.	/_	/	

		BLOOD SUGAR RESULTS*							
		Breakfast				Din	Dinner		Night
	MONDAY	BEFORE	AFTER	BEFORE	AFTER	BEFORE	AFTER	Time	Time
Medicine type and dose ▼	units	Time	Time	Time	Time	Time			
ty OSE									
d e.		mg/dL	ma/dl	ma/dl	ma/dl	mg/dL	ma/dl	mg/dL	mg/dL
Sign	Carb intake	IIIg/uL	IIIg/uL	IIIg/uL	IIIg/uL	IIIg/uL	IIIg/uL	Hig/uL	IIIg/uL
Je (Mealtime insulin dose	Rroakfact	unite	□Lunch	unite	□ Dinner	unite		
_	Other >	DIGANIASI	uiiis	Lulion	uiiis		uiiits		
	Other	Brea	kfast	Lur	nch	Din	ner	Redtime	Night
	TUESDAY							Time	Time
e N	units		Time	Time	Time	Time	Time	Timo	Timo
t Se									
9									
<u>.</u> <u>D</u> <u>D</u>		mg/dL	mg/dL	mg/dL	mg/dL	mg/dL	mg/dL	mg/dL	mg/dL
Medicine type and dose ▼	Carb intake ►								
2	Mealtime insulin dose	☐ Breakfast	units	Lunch	units	☐ Dinner	units		
	Other►	Dura			1-	D:		D. It'	Ni ulat
	MEDNICOAV	Breal	kfast	LUI	1CN	DEFORE	ner	Bedtime	
ا ل	WEDNESDAY units	Time		Time	Time	Time	Time	Time	Time
S S	- Cirio	TITIO	Time	Tillio	Tillio	Time	TITIE		
Medicine type and dose ▼									
<u>a</u>		mg/dL	mg/dL	mg/dL	mg/dL	mg/dL	mg/dL	mg/dL	mg/dL
ed	Carb intake ►								
Σ	Mealtime insulin dose >	☐ Breakfast	units	Lunch	units	☐ Dinner	units		
	Other ►								
		Brea	kfast	Lur	<u>nch</u>	Din	ner	Bedtime	Night
a) .	THURSDAY units	BEFORE	AFTER	BEFORE	AFTER			Time	Time
o be	units	Time	Time	Time	Time	Time	Time		
e t									
9		mg/dL	mg/dL	mg/dL	mg/dL	mg/dL	mg/dL	mg/dL	mg/dL
Medicine type and dose ▼	Carb intake >	, and the second		, and the second					
Š	Mealtime insulin dose	☐ Breakfast _	units	Lunch	units	☐ Dinner	units		
	Other ►								
		Brea	kfast	Lur	nch	Din	ner	Bedtime	Night
.	FRIDAY	BEFORE	AFTER					Time	Time
ledicine type and dose ▼	units	Time	Time	Time	Time	Time	Time		
e t)									
d d		ma/dl	mg/dL	ma/dl	ma/dl	ma/dl	ma/dl	mg/dL	mg/dL
a dic	Carb intake ►	g,	9,	g,	9,	9,	9	g/ ==	
\mathbb{R}	Mealtime insulin dose	☐ Breakfast	units	Lunch	units	☐ Dinner	units		
	Other >								
		Brea	kfast	Lur	nch	Din	ner	Bedtime	Night
	SATURDAY	BEFORE	AFTER	BEFORE		BEFORE		Time	Time
be	units	Time	Time	Time	Time	Time	Time		
ty OSE									
<u>a</u> <u>a</u> <u>a</u>		mg/dL	mg/dL	mg/dL	mg/dL	mg/dL	mg/dL	mg/dL	mg/dL
Medicine type and dose ▼	Carb intake ▶	IIIg/uL	IIIg/uL	IIIg/uL	IIIg/uL	IIIg/uL	IIIg/uL	IIIg/uL	IIIg/uL
Ş,	Mealtime insulin dose	☐ Breakfast	units	Lunch	unite	☐ Dinner	units		
_	Other	Dicariast	units	Lunon	uiiio		units		
	Other	Brea	kfast	Lur	nch	Din	ner	Bedtime	Night
oc •	SUNDAY		AFTER		AFTER	BEFORE	AFTER	Time	Time
	units	Time	Time	Time	Time	Time	Time		
t Se									
9									
n di		mg/dL	mg/dL	mg/dL	mg/dL	mg/dL	mg/dL	mg/dL	mg/dL
Medicine type and dose ▼	Carb intake	Description 1			. 9	□ Diama			
2	Mealtime insulin dose	☐ Breaktast _	units	LI LUHCH	units	☐ Dinner	units		

n